

Combinatorial Matrices' Kallista symbol: Projective, Spectral, isotropic and multiplicative

E. Anderson*

Abstract

Combinatorial matrices have recently been studied by Ford and the Author. We here add a fourth kind of Ford symbol, which carries both Projective and Spectral connotations. And unlike the previous three, manages to have the very simplest multiplicative property. Revealing that Combinatorial matrices' eigenvalues combine in the simplest possible way under products and sums: an uncommon feature among square matrices. $K \times K$ Combinatorial matrices support not only K -square eigenexpansions but also 2-square eigenexpansions. This is underpinned by Combinatorial matrices mostly consisting of an isotropic block. A new style of proof for each K 's Combinatorial matrices commuting with each other is also included.

* Dr.E.Anderson.Maths.Physics *at* protonmail.com . Institute for the Theory of STEM.

Date stamp: v1 03-10-2025, v2 04-11-2025 Copyright of Dr E. Anderson.

1 Introduction

Definition 1 A *Combinatorial matrix* [1] is a $K \times K$ matrix of the following form.

$$C := \begin{pmatrix} x+y & x & \dots & x \\ x & & & \vdots \\ \vdots & & & x \\ x & \dots & x & x+y \end{pmatrix} = x\mathbb{1} + y\mathbb{I} =: (x+y, y)_K. \quad (1)$$

Remark 1 While Combinatorics often involves $x, y \in \mathbb{Z}$ or \mathbb{N} , Flat Geometry applications [8, 9, 10] extend interest to \mathbb{R} . Here \mathbb{I} is the identity matrix and $\mathbb{1}$ is the matrix of 1s. Also $(\cdot, \cdot)_K$ is the Ford symbol of the zeroth kind [12, 13], a truer name for which is *irreducible symbol*. With reference to the trace-tracefree basis, with basis matrices \mathbb{I} and \mathbb{T} : 1's off-diagonal and 0's on-diagonal. Whenever we work with a fixed K , we simplify this notation to (\cdot, \cdot) .

Remark 2 For a fixed K , the totality of these constitute the arena $\mathbf{CM}_K(2)$: a 2-d vector space [12].

Remark 3 Combinatorial matrices were expanded relative to various other bases in [10, 12]. We now consider a further such, which accumulates various interesting properties.

Some previous bases involved \mathbb{I} . We consider this again, now however viewing

$$K^{-1}\mathbb{I} = (1, 1)$$

in the following further way. As the dyad of normalized equal-entries vectors

$$\bar{\mathbf{n}}\mathbf{n}.$$

Which is furthermore an orthogonal projector

$$\bar{\mathbf{P}}_{\perp} \quad (2)$$

onto the 1-d total-sum direction space [13]

$$\mathbf{Dir}(1). \quad (3)$$

Whose complement is k -dimensional difference space [13]

$$\mathbf{Dir}(k). \quad (4)$$

With corresponding orthogonal projector

$$\bar{\mathbf{P}}_{\perp\perp} = \bar{\mathbf{P}} = \mathbb{I} - \bar{\mathbf{n}}\mathbf{n}. \quad (5)$$

For Combinatorial matrices are blessed with [13] an at least k -fold degenerate eigenvalue. Where

$$k := K - 1.$$

Resting upon at least an $O(k)$ -symmetry isotropy being forced [13]. In the generic case, the corresponding eigensubspace is difference space. The only alternative to this is conflation with the remaining eigenvalue. Yielding an all-encompassing K -fold degenerate eigenspace. Now resting upon the whole matrix being $O(K)$ -symmetry isotropic [13].

In the N -body problem context, (3) is the centre of mass (CoM) position label. While (5) is the CoM-removing projector And (4) takes the form of relative label space

$$\mathfrak{Rel}(n) = \mathbb{R}^n :$$

the space of linearly-independent (LI) separation vector labels. Labels first enter 1, 2, ... to the points. Leading to the separations being labelled by their pair of boundary points, e.g. 1 2. CoM position carries both a spatial vector index and a point label index. So does the relative space of LI separations [18]. But which are only the label part plays an active role: the full version of everything just tensors with the spatial identity matrix [18].

In this context, \mathbf{P} is furthermore numerically (if not Physical-dimensionally [18]) the Lagrange matrix [16]. The overarching theme is that upon translating to the CoM frame, all remaining degrees of freedom are relative separations. With the Lagrange matrix arising by [18] extremizing the arbitrary-translation correction to the inertia quadric with respect to its translation variable. This is how translations, CoMs and separations are inter-related. For the general Combinatorial matrix, upon translating to the *total sum*, all remaining freedom lies correspondingly in *difference variables* [13].

Remark 4 The current Article's incipient idea is to take \mathbf{P}_\perp and

$$\mathbf{P} = K^{-1}(k, -1) \quad (6)$$

as our basis. Which is a priori motivated by its being privileged by both of its basis matrices being projectors. While the below results build up additional a posteriori motivations.

2 Eigentheory results

Lemma 1

$$(x + y, x)\mathbf{P} = y\mathbf{P} = \lambda\mathbf{P} =: \lambda_k\mathbf{P}. \quad (7)$$

$$(x + y, x)\mathbf{P}_\perp = (Kx + y)\mathbf{P}_\perp = \lambda_\perp\mathbf{P}_\perp =: \lambda_1\mathbf{P}_\perp. \quad (8)$$

Where λ is the k -fold degenerate eigenvalue λ_k : with algebraic multiplicity k [13]. While λ_\perp is the complementary eigenspace's lone eigenvalue λ_1 : with algebraic multiplicity 1 .

Corollary 1

$$(x + y, x) = \lambda\mathbf{P} + \lambda_\perp\mathbf{P}_\perp. \quad (9)$$

Naming Remark 1 This prompts introducing a fourth 'Ford symbol': the *Projective symbol*

$$\{\lambda, \lambda_\perp\}_K. \quad (10)$$

Where the first entry is the \mathbf{P} component and the second the \mathbf{P}_\perp component. But these entries are furthermore the spectrum of eigenvalues (without multiplicities). And so another truer name for it is *Spectral symbol*. That a pair of eigenvalues covers the whole spectrum in this sense is underpinned by the above observation that Combinatorial matrices have an at least k -fold degenerate eigenvalue [13]. Which rests in turn on an at least $O(k)$ -symmetry isotropy. Whence a third name: *isotropy symbol* with reference to an at-most codimension-1 isotropy.

Lemma 2

$$\{p\lambda_1, p\lambda_{2\perp}\} = p\{\lambda_1, \lambda_{2\perp}\}. \quad (11)$$

$$\{\lambda_1, \lambda_{1\perp}\} + \{\lambda_2, \lambda_{2\perp}\} = \{\lambda_1 + \lambda_2, \lambda_{1\perp} + \lambda_{2\perp}\}. \quad (12)$$

$$\{\lambda_1, \lambda_{1\perp}\}\{\lambda_2, \lambda_{2\perp}\} = \{\lambda_1\lambda_2, \lambda_{1\perp}\lambda_{2\perp}\}. \quad (13)$$

Remark 1 So passing to using our symbol preserves the simplest linearity properties, while attaining the simplest product formula. Nor is it at all usual for the eigenvalues of the product of 2 matrices

to be the product of the eigenvalues of the 2 matrices. And yet by the above this holds for any 2 Combinatorial matrices of the same size... And similarly with ‘sum’ in place of ‘product’!

Naming Remark 2 Let us celebrate by, firstly, placing a fourth name on the Combinatorial matrix symbol $\{, \}$: *multiplicative symbol*. Secondly, by pointing out that at present this symbol ‘conceptually outnumbers’ our irreducible symbol $(,)$: by 4 to 1 . I.e. it is known to have multiplicative, Projective, Spectral and isotropic significance. To its rival carrying Representation-Theoretic significance. By the above confluence of properties from across Mathematics, we call $\{, \}$ the Combinatorial matrix *Kallista symbol*. And thirdly by extending to the following.

Corollary 2

$$\sum_{i=1}^R \{ \lambda_i, \lambda_{i \perp} \} = \left\{ \sum_{i=1}^R \lambda_i, \sum_{i=1}^R \lambda_{i \perp} \right\}, \quad \prod_{i=1}^R \{ \lambda_i, \lambda_{i \perp} \} = \left\{ \prod_{i=1}^R \lambda_i, \prod_{i=1}^R \lambda_{i \perp} \right\}. \quad (14)$$

♠ ♦ ♣

Remark 2 Next working with the quadratic form version – acting on an input vector of numbers \mathbf{K} , we readily obtain the following results. In the N -body problem case, it turns out to be quite useful [9, 18] to take $\mathbf{K} = \mathbf{S}$: the vector of side-lengths squared. These are 1 dependency away from being LI. But there are also K of them, while we are only expecting k differences. So everything checks out at the level of well-determinedness.

Proposition 1 2-squares expansion.

$$\|\mathbf{K}\|_{\{ \lambda, \lambda_{\perp} \}}^2 = \lambda_{\perp} \|\mathbf{K}\|_{\mathbf{P}_{\perp}}^2 + \lambda \|\mathbf{K}\|_{\mathbf{P}}^2 = \lambda_{\perp} K^{-1} U^2 + \lambda \|\mathbf{K}\|_{\mathbf{P}}^2. \quad (15)$$

Proposition 2 The K -squares expansion in $\mathbf{K}(K)$ network choice of basis is as follows.

$$\|\mathbf{K}\|_{\{ \lambda, \lambda_{\perp} \}}^2 = \lambda_{\perp} K^{-1} U^2 + \lambda \sum_{q=1}^k q^{-1} Q^{-1} \left(\sum_{p=q}^k K_p - q K_Q \right)^2. \quad (16)$$

Remark 3 In the N -body problem context, firstly the unit-normalized total sum of the objects U is realized by a quantity proportional to R : the square of the *radius of gyration*. Secondly, the first distinct K network is called the Jacobi- K [3, 16, 18] for the 4-body problem. This corresponds to [6, 11] the straight-3-path unlabelled rooted binary tree (URBT) [2]. The subsequent K networks are the corresponding straight-path URBTs for each larger path. [13] showed that these bases remain meaningful for arbitrary Combinatorial matrices. Thirdly, the above codimension-1 isotropy corresponds in the N -body setting to the $O(n)$ group of *internal rotations* alias *democracy transformations* [4].

Corollary 3

$$\|\boldsymbol{\kappa}\|_{\{ \lambda, \lambda_{\perp} \}}^2 = \lambda_{\perp} K^{-1} + \lambda \|\boldsymbol{\kappa}\|_{\mathbf{P}}^2 = \lambda_{\perp} K^{-1} + \lambda \sum_{q=1}^k q^{-1} Q^{-1} \left\| \sum_{p=q}^k \boldsymbol{\kappa}_p - q \boldsymbol{\kappa}_Q \right\|^2. \quad (17)$$

Remark 4 Where we have introduced the ratio variables

$$\boldsymbol{\kappa} := U^{-1} \mathbf{K}. \quad (18)$$

As enabled by the above at most codimension-1 isotropy.

Remark 5 The K -squares expansion uses the whole eigenspectrum – with algebraic multiplicity – based on

$$\mathfrak{M}_K = \text{Im}_{\mathbf{P}_{\perp}}(\mathfrak{M}_K) \oplus \text{Im}_{\mathbf{P}}(\mathfrak{M}_K) = \mathfrak{D}\text{ir}(1) \oplus \mathfrak{D}\text{ir}(k).$$

For \mathfrak{M}_K the space of $K \times K$ matrices. In contrast, 2-squares expansion uses that the space of Combinatorial matrices is that

$$\mathfrak{CM}_K(2) = \mathfrak{D}\text{ir}(1) \oplus \mathfrak{D}\text{ir}(1).$$

3 Examples

Examples 1-7: in our brackets notation. The CoM-removing projector itself is

$$\mathbf{P} = \{1, 0\}.$$

Its complement is

$$\mathbf{P}_\perp = \{0, 1\}.$$

The zero matrix is

$$\mathbf{0} = \{0, 0\}.$$

And the identity matrix is

$$\mathbf{I} = \{1, 1\}.$$

The previous two are both examples of isotropic matrices, the general case of which is

$$r\{1, 1\}, \quad r \in \mathbb{R}.$$

The fundamental 2-simplex matrix, shared by the triangle inequality, the cosine rule, and Heron's formula, [7, 8, 9, 10] is

$$\mathbf{F} = \{-2, 1\}.$$

The Apollonius involutor [7, 9, 10] is

$$\mathbf{J} = \frac{1}{3}\{-1, 1\}.$$

Example 8 Of sums of K squares. In the $K = N = 3$ -body problem context, Proposition 2 returns $Aniso^2 + Anelp^2$ for its last 2 squares [7, 8, 9, 18]. Standing for *anisoscelesness*: departure from isoscelesness. And departure from being in equilateral proportion, with reference to the base to median ratio. In terms of side-lengths² A, B, C ,

$$Aniso = \frac{A - B}{\sqrt{2}}, \quad Anelp = \frac{A + B - 2C}{\sqrt{6}}.$$

Generalizing from sides-lengths² in the 3-body problem to differences of primary objects in Combinatorics, we write U in place of R . Not *Aniso* but *Ind*, in the sense of induced from a $K = 2$ difference. And not *Anelp* but *Comp*, in the sense of orthonormal complement.

Proposition 2 recovers as a subcase

$$\|\mathbf{K}\|_{(x+y, x)}^2 = \frac{3x + y}{3} U^2 + \frac{y}{2} \left[(A - B)^2 + \frac{(A + B - 2C)^2}{3} \right]. \quad (19)$$

Which can also be written in the following generalized Euler 3-cycle form.

$$\begin{aligned} \|\mathbf{K}\|_{(x+y, x)}^2 &= \frac{3x + y}{3} \sum_{3\text{-cycles}} A(A + 2B) + \frac{2y}{3} \sum_{3\text{-cycles}} A(A - B) \\ &= \sum_{3\text{-cycles}} A[(x + y)A + 2xB]. \end{aligned} \quad (20)$$

Setting x and y to match \mathbf{F}, \mathbf{J} and \mathbf{P} in turn recovers some previous results from [7, 10]. In particular, since \mathbf{P} is itself the projector, the eigenvalue λ_\perp corresponding to \mathbf{P}_\perp is 0. And so the $U = R$ contribution to \mathbf{P}_\perp drops out entirely.

4 Algebras

**Combinatorial matrix
projectors'
commuting monoid P**

•	1	P	P_\perp	0
1	1	P	P_\perp	0
P	P	P	0	0
P_\perp	P_\perp	0	P_\perp	0
0	0	0	0	0

© 2025 Dr E. Anderson

Figure 1:

Proposition 3 Under multiplication, P and P_\perp alongside the identity 1 and zero 0 form the commutative monoid [5] whose times table is in Fig 1.

Proposition 4 The above four inputs also form a zero-commutator algebra.

Remark 1 Proposition 3's times table follows from just the identity property, the projector property and projector complementarity. Proposition 3's inputs reflect that, on the one hand, we can freely append an identity element. On the other hand, we are forced to include the zero since we discover it as the product of our two projectors.

Remark 2 For Proposition 4, the only nontrivial bracket to check is

$$[\bar{P}, \bar{P}_\perp] = \bar{P} \cdot \bar{P}_\perp - \bar{P}_\perp \cdot \bar{P} = \bar{0} - \bar{0} = \bar{0}.$$

Where step 1 is by the definition of commutator. And step 2 makes 2 uses of complementarity.

Remark 3 The current Section is the generalized Combinatorial matrices' counterpart of the triangle (or more generally 2-simplex) algebras presented in [10]. And of various quadrilateral algebras presented in [14, 15, 17].

5 Conclusion

Remark 1 A first motivation for our Kallista symbol for Combinatorial matrices is that it corresponds to the Projective choice of basis. In the current Article, this appeared as our a priori reason to entertain this symbol. A posteriori, we found 3 further motivations to strengthen this.

Remark 2 Using our new basis, the product rule (13) and the commutative monoid concurrently manage to take particularly simple forms. In contrast to the various basis choices in [10], the current Article's commutative monoid's is distinctive by its times table necessarily involving the zero. That compatible Combinatorial matrices always commute was recently pointed out by Ford [12]. Proposition 4 however provides an alternative style of proof for this! I.e. every Combinatorial matrix can be written as the weighted sum of a complementary pair of projectors. Which commute with each other, and so all compatible sized Combinatorial matrices commute with each other. This paragraph lays out the multiplicative motivation for the Kallista symbol.

Remark 3 A third motivation is Spectral: our basis displays the eigenvalues as its components. And clarifies that these combine particularly simply for Combinatorial matrices under both addition and multiplication. General square matrices' eigenvalues do not have these properties! By which our basis displays a 2-term eigenexpansion. This is rendered possible by all Combinatorial matrices having an eigenvalue that is at least k -fold degenerate by symmetry. Due to the underlying at-least $O(k)$ -symmetry – and thus at most codimension-1 – isotropy: a fourth Group-Theoretic motivation.

Acknowledgments I thank A. Ford and K. Everard for discussions about Combinatorial matrices and Linear Algebra. And S. Sánchez for past discussions about Geometry, Combinatorics, Applied Topology, and the Mathematical citizens of Kallista more generally. This work follows on from the “Linear Algebra of Quadrilaterals” Summer School 2024 at the Institute for the Theory of STEM.

References

- [1] D.E. Knuth, *The Art of Computer Programming 1 Fundamental Algorithms* (Addison-Wesley, 1968).
- [2] **3 Sorting and Searching** (Addison-Wesley, 1975).
- [3] R.G. Littlejohn and M. Reinsch, “Internal or Shape Coordinates in the N-Body Problem”, *Phys. Rev. A* **52** 2035 (1995).
- [4] “Gauge Fields in the Separation of Rotations and Internal Motions in the N-Body Problem”, *Rev. Mod. Phys.* **69** 213 (1997).
- [5] P.M. Cohn, *Basic Algebra. Groups, Rings and Fields* (Springer, London 2003).
- [6] D. Schmidt, “Central Configurations and Relative Equilibria for the N-Body Problem”, in *Classical and Celestial Mechanics* ed. H Cabral and F. Diacu (P.U.P. , Princeton NJ 2002).
- [7] E. Anderson, “Two New Perspectives on Heron’s Formula”, *Ann. Inst. Theory STEM* **1** 1 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> . arXiv:1712.01441; Updated preprint: <https://wordpress.com/page/conceptsofshape.space/1244> (2024).
- [8] “The Fundamental Triangle Matrix”, *Ann. Inst. Theory STEM* **1** 2 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> . Preprint: <https://wordpress.com/page/conceptsofshape.space/1306> (2024).
- [9] “The Fundamental Triangle Matrix Commutes with the Lagrange and Apollonius Matrices. With implications for deriving Kendall and Hopf Theorems.” *Ann. Inst. Theory STEM* **1** 3 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> . Preprint: <https://wordpress.com/page/conceptsofshape.space/1308> (2024).
- [10] “Only 2 of the Fundamental Triangle, Lagrange and Apollonius Matrices are Independent. With ensuing Algebras, Irreducibles and Splits” *Ann. Inst. Theory STEM* **1** 4 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> . Preprint: <https://wordpress.com/page/conceptsofshape.space/1310> (2024).
- [11] E. Anderson and A. Ford, “Graph and Order Theory of N-Body Problem’s Eigenclustering Networks”, accepted by Period. Updating Rev. App. Top. Preprint: <https://wordpress.com/page/conceptsofshape.space/2090> (2024).
- [12] A. Ford, “Triangle, but not most Quadrilateral, Matrices are Combinatorial” *Ann. Inst. Theory STEM* **1** 5 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> .
- [13] E. Anderson, “Eigentheory of Combinatorial Matrices: a More General Setting for the CoM-Relative Split”, *Ann. Inst. Theory STEM* **1** 6 (2025), <https://institute-theory-stem.org/annals-of-the-institute-of-stem/> . Preprint: <https://wordpress.com/page/conceptsofshape.space/2280> (2024).
- [14] “4-Body Problem: Ptolemy–Lagrange Algebra”, Preprint: <https://wordpress.com/page/conceptsofshape.space/2121> (2024).
- [15] E. Anderson and K. Everard, “Linear Algebra of Cyclic Quadrilaterals: Ptolemy, Diagonal, Area, and Circumradius Formulae”, Preprint: <https://wordpress.com/page/conceptsofshape.space/1971> (2024) .
- [16] R. Montgomery, *Four Open Questions for the N-Body Problem* (C.U.P., Cambridge 2025).
- [17] E. Anderson, “Convex Quadrilaterals’ Ptolemy–Lagrange–Bretschneider Algebras”, Preprint: <https://wordpress.com/page/conceptsofshape.space/2278> (2025).
- [18] *The Structure of Flat Geometry*, in some places called additionally *Widely-Applicable Mathematics. A. Improving understanding of everything by dual-wielding Combinatorics and Linear Algebra.* **3**, forthcoming 2026.